

Effective string action for the $U(1) \times U(1)$ dual Ginzburg-Landau theory beyond the London limit

Yoshiaki Koma^{a,b}, Miho Koma^{c,b}, Dietmar Ebert^d, Hiroshi Toki^c

^a *Institute for Theoretical Physics, Kanazawa University,
Kanazawa, Ishikawa 920-1192, Japan*

^b *Max-Planck-Institut für Physik, Föhringer Ring 6, D-80805 München, Germany*

^c *Research Center for Nuclear Physics, Osaka University,
Mihogaoka 10-1, Ibaraki, Osaka 567-0047, Japan*

^d *Institut für Physik, Humboldt Universität zu Berlin, D-10115 Berlin, Germany*

Abstract

The effective string action of the color-electric flux tube in the $U(1) \times U(1)$ dual Ginzburg-Landau (DGL) theory is studied by performing a path-integral analysis by taking into account the finite thickness of the flux tube. The DGL theory, corresponding to the low-energy effective theory of Abelian-projected $SU(3)$ gluodynamics, can be expressed as a $[U(1)]^3$ dual Abelian Higgs (DAH) model with a certain constraint in the Weyl symmetric formulation. This formulation allows us to adopt quite similar path-integral techniques as in the $U(1)$ DAH model, and therefore, the resulting effective string action in the $U(1) \times U(1)$ DGL theory has also quite a similar structure except the number of color degrees of freedom. A modified Yukawa interaction appears as a boundary contribution, which is completely due to the finite thickness of the flux tube, and is reduced into the ordinary Yukawa interaction in the deep type-II (London) Limit.

Key words: Effective string action, dual Ginzburg-Landau theory, confinement
PACS: 12.38.Aw, 12.38.Lg

Email addresses: ykoma@mppmu.mpg.de (Yoshiaki Koma), mkoma@mppmu.mpg.de (Miho Koma), debert@physik.hu-berlin.de (Dietmar Ebert), toki@rcnp.osaka-u.ac.jp (Hiroshi Toki).

1 Introduction

The dual superconducting picture of the QCD vacuum [1,2] provides an intuitive scenario on the quark confinement mechanism. The formation of the color-electric flux tube can be understood analogously to that of the magnetic Abrikosov vortex [3,4,5] in an ordinary superconductor via the Meissner effect with the electric Cooper-pair condensation. In fact, in the QCD vacuum, color-magnetic monopoles are condensed, and then, the color-electric field associated with quarks is squeezed into an almost one-dimensional object forming the color-electric flux tube due to the dual Meissner effect (the dual Higgs mechanism). The linear rising potential naturally appears between the quark and the antiquark, which explains why a quark cannot be isolated as a free particle.

Remarkably, color-magnetic monopole condensation is observed by Monte-Carlo simulations of lattice gluodynamics in the so-called maximally Abelian (MA) gauge [6,7,8]. In the MA gauge, Abelian dominance [9,10] for the non-perturbative quantities, such as the string tension [11,12] and chiral condensate [13], is also observed. Moreover, it is suggested that monopole configurations lead to large contributions to the string tension [14] and chiral condensate [15]. These results indicate that Abelian components of the gluon field and monopoles become relevant degrees of freedom to describe the low-energy (long-range) properties of the vacuum.

These numerical results have justified a posteriori a low-energy effective theory of the gluodynamics composed of Abelian degrees of freedom (Abelian-projected gluodynamics [16]) with monopole condensation. The resulting effective action is known to have the form of the Ginzburg-Landau (GL) theory which in its original version would effectively describe bulk properties of the ordinary superconductor. Since now the role of electric and magnetic properties is interchanged, this is called the dual Ginzburg-Landau (DGL) theory [17,18,19,20]. The DGL Lagrangian is composed of the dual gluon field interacting with the monopole field, and in fact, the color-electric flux-tube configuration is obtained as the classical solution when the test electric charges are put into the vacuum. The thickness of the flux tube is controlled by the inverse masses of the dual gauge boson and the monopole, that is, the penetration depth m_B^{-1} and the coherence length m_χ^{-1} . The latter length characterizes the so-called core size of the flux tube.

Although long-distance aspects are dominated by Abelian-projected degrees of freedom, the DGL theory may not be a complete substitute for all aspects of low-energy gluodynamics, since there would be an interface in the real theory with short range physics which is taken care of by charged vector and ghost fields. Such effects would play an important role for the description of

a dynamical charge density around test charges. Nevertheless, it is interesting to remind that the Abelian-projected flux-tube profiles have also been studied within the lattice gluodynamics in the MA gauge [21,22,23,24,25,26], which have shown quite similar profiles as the classical flux-tube solution in the DGL theory. Therefore it will make sense to regard the DGL theory as a starting point of further investigations of the hadronic flux tube [27,28,29,30] with the pointlike approximation of color-electric test charges.¹

From the Abelian-projected $SU(2)$ gluodynamics one can construct the $U(1)$ DGL theory (often called the $U(1)$ dual Abelian Higgs (DAH) model), where the $U(1)$ dual symmetry corresponds to the Cartan subgroup of $SU(2)$. One can analyze the physical essence of the dual superconducting vacuum and, at the same time, the structure of the flux tube. For the description of the dynamics of the flux tube, it makes sense to use a framework which directly deals with the string-like properties of the flux tube. Such framework is the *string representation* or *effective string action approach* of the DGL theory, which can be obtained by performing a path-integral analysis. In the $U(1)$ case, based for simplicity, on the London limit approximation $m_\chi \rightarrow \infty$, the string representation was derived in Refs. [31,32,33,34]. The London limit approximation corresponds to the neglect of the core size of the flux tube. The resulting leading action was shown to be the Nambu-Goto (NG) action with the rigidity term [35,36]. The string representation of the $U(1) \times U(1)$ DGL theory corresponding to the Abelian-projected $SU(3)$ gluodynamics was also studied using this approximation [37,38].

Recently, we have considered the derivation of the string representation of the $U(1)$ DGL theory beyond the London limit, *i.e.* we have taken into account the finite thickness of the flux-tube core by keeping the mass of the monopole finite [39]. The used path-integral analysis combined with differential form techniques turned out to provide a very compact and transparent method for studying these problems including, in particular, also boundary terms related to the non-confining part of the potential. As a main point, the finite core corrections to the effective string action have been derived.

In this paper, we are going to generalize the above work by investigating the $U(1) \times U(1)$ DGL theory beyond the London limit. Note that two of us have shown that the Weyl symmetric formulation reduces the $U(1) \times U(1)$ DGL theory into the sum of three types of the $U(1)$ DAH model, which was quite useful to simplify the treatment of the DGL theory and the investigation of the flux-tube solution [29,30]. Therefore, in order to derive the effective string action, we pay here special attention to the Weyl symmetric formulation when using again path-integral techniques in differential form notation.

¹ The resulting DGL description is thus, in particular, justified for the string away from the charges and their dynamical charge clouds.

This paper is organized as follows. In Sec. 2 we introduce the Weyl symmetric formulation of the DGL theory in differential form notation. In Sec. 3 the role of the open Dirac string with color-electric charge at its ends for the violation of the dual Bianchi identity is considered. In Sec. 4 the functional path integration in differential form notation is performed which requires special attention to gauge conditions and the Weyl symmetry constraint. This leads to the effective string action, where the finite thickness of the core of the flux tube is taken into account. Finally, Sec. 5 contains the summary.

2 Weyl symmetric formulation of the DGL theory

The Weyl symmetric expression of the DGL action in the differential form is given by

$$S_{\text{DGL}} = \sum_{i=1}^3 \left[\frac{\beta_g}{2} F_i^2 + ((d - iB_i)\chi_i^*, (d + iB_i)\chi_i) + \lambda(|\chi_i|^2 - v^2)^2 \right], \quad (1)$$

where the ingredients of the DGL theory are three types of the dual gauge field B_i ($\sum_{i=1}^3 B_i = 0$, $i = 1, 2, 3$), and the complex-scalar monopole field $\chi_i = \phi_i \exp(i\eta_i)$ ($\phi_i, \eta_i \in \mathfrak{R}$, $i = 1, 2, 3$). In the presence of external quark sources in the vacuum describing a static quark-antiquark ($q\bar{q}$) system, the dual field strength F_i is expressed as

$$F_i = dB_i - 2\pi * \Sigma_i^{(m)}, \quad (2)$$

with

$$* \Sigma_i^{(m)} \equiv \sum_{j=1}^3 m_{ij} * \Sigma_j^{(e)}, \quad (3)$$

Here $\Sigma_i^{(m)}$ ($i = 1, 2, 3$) denotes the “color-magnetic” (Weyl symmetric) representation of the color-electric Dirac string $\Sigma_j^{(e)}$ ($j = 1, 2, 3$) which describes the *open* color-electric Dirac string [30]. Let us briefly summarize the definitions of operators in differential forms and useful formulae in Table 1, and quote the ingredients of the DGL theory in Table 2. The factor $2\pi m_{ij}$ in front of the color-electric Dirac string appears as a result of the extended Dirac quantization condition between the color-magnetic charges $\vec{Q}_i^{(m)} \equiv g\vec{e}_i$ ($i = 1, 2, 3$) and color-electric charges $\vec{Q}_j^{(e)} \equiv e\vec{w}_j$ ($j = 1, 2, 3$):

$$\vec{Q}_i^{(m)} \cdot \vec{Q}_j^{(e)} = g\vec{e}_i \cdot e\vec{w}_j = 2\pi m_{ij} \quad (eg = 4\pi), \quad (4)$$

Table 1

Definitions in differential forms in four-dimensional Euclidean space-time.

r -form ($0 \leq r \leq 4$)	ω	$\omega \equiv \frac{1}{r!} \omega_{\mu_1 \dots \mu_r} dx_{\mu_1} \wedge \dots \wedge dx_{\mu_r}$
exterior derivative	d	r -form $\mapsto (r+1)$ -form
Hodge star	$*$	r -form $\mapsto (4-r)$ -form
	$**$	multiply by a factor $(-1)^r$ for an r -form
codifferential	$\delta \equiv - * d *$	r -form $\mapsto (r-1)$ -form
Laplacian	$\Delta \equiv d\delta + \delta d$	r -form $\mapsto r$ -form
Inner product	$(\omega, \eta) \equiv \int \omega \wedge * \eta$	$(\omega, \eta) = \frac{1}{r!} \int d^4x \omega_{\mu_1 \dots \mu_r} \eta_{\mu_1 \dots \mu_r} \quad (\omega, \eta \in r\text{-form})$
		$(\omega)^2 \equiv (\omega, \omega)$

Table 2

Ingredients of the DGL theory in differential-form notation. The subscripts $i = 1, 2, 3$ denote the label of color in the color-magnetic (Weyl symmetric) representation.

dual gauge field	1-form	$B_i \equiv B_{i\mu} dx_\mu$
monopole field	0-form	$\chi_i \equiv \phi_i \exp(i\eta_i)$
electric Dirac string	2-form	$\Sigma_i^{(m)} \equiv \frac{1}{2} \Sigma_{i\mu\nu}^{(m)} dx_\mu \wedge dx_\nu$
electric current	1-form	$j_i^{(m)} \equiv j_{i\mu}^{(m)} dx_\mu$

where $\vec{\epsilon}_i$ and $\vec{w}_j$ are the root and weight vectors of the SU(3) Lie algebra, respectively. Explicitly, these vectors are defined as $\vec{\epsilon}_1 = (-1/2, \sqrt{3}/2)$, $\vec{\epsilon}_2 = (-1/2, -\sqrt{3}/2)$, $\vec{\epsilon}_3 = (1, 0)$ and $\vec{w}_1 = (1/2, \sqrt{3}/6)$, $\vec{w}_2 = (-1/2, \sqrt{3}/6)$, $\vec{w}_3 = (0, -1/\sqrt{3})$. The labels $j = 1, 2, 3$ describe, for instance, red (R), blue (B), and green (G) color-electric charges, respectively. The matrix

$$m_{ij} \equiv 2\vec{\epsilon}_i \cdot \vec{w}_j = \sum_{k=1}^3 \varepsilon_{ijk} \quad (5)$$

takes the integer values ± 1 or 0 . Based on these definitions of charges, the R - $\bar{R}$ system is defined by $\Sigma_{j=1}^{(e)} = \Sigma \neq 0$, $\Sigma_{j=2}^{(e)} = 0$, and $\Sigma_{j=3}^{(e)} = 0$, which is also expressed as $\Sigma_{i=1}^{(m)} = 0$, $\Sigma_{i=2}^{(m)} = -\Sigma$, and $\Sigma_{i=3}^{(m)} = \Sigma$ in the color-magnetic representation. Clearly, the color-electric Dirac string in the color-magnetic representation satisfies the relation $\sum_{i=1}^3 \Sigma_i^{(m)} = 0$ (see, Table 3).

The physics described by the DGL theory is characterized by two mass scales, the masses of the dual gauge boson $m_B = \sqrt{2/\beta_g} v = \sqrt{3} g v$ and of the monopole $m_\chi = 2\sqrt{\lambda} v$, where $\beta_g = 2/(3g^2)$ is the dual gauge coupling and λ the self-coupling of the monopole field. Recall that their inverses correspond to the penetration depth and the coherence length, respectively. We can further

Table 3

The color-electric Dirac string structures in the color-electric and color-magnetic representations for the $q-\bar{q}$ system, where Σ corresponds to the world sheet of one string singularity.

	color-electric rep.			color-magnetic rep.		
	$\Sigma_{j=1}^{(e)}$	$\Sigma_{j=2}^{(e)}$	$\Sigma_{j=3}^{(e)}$	$\Sigma_{i=1}^{(m)}$	$\Sigma_{i=2}^{(m)}$	$\Sigma_{i=3}^{(m)}$
$R-\bar{R}$	Σ	0	0	0	$-\Sigma$	Σ
$B-\bar{B}$	0	Σ	0	Σ	0	$-\Sigma$
$G-\bar{G}$	0	0	Σ	$-\Sigma$	Σ	0

define the Ginzburg-Landau (GL) parameter κ as

$$\kappa \equiv m_\chi/m_B, \quad (6)$$

which controls the vacuum property of dual superconductivity, in particular, whether the vacuum belongs to type-I ($\kappa < 1$) or type-II ($\kappa > 1$) or to their border ($\kappa = 1$) .

3 Open string singularity and the violation of the dual Bianchi identity

When we consider the open string system, such as the $R-\bar{R}$ system, it is important to pay attention to the color-electric Dirac string singularity in the dual field strength F_i , since it is directly related to the violation of the dual Bianchi identity:

$$dF_i = d^2 B_i - 2\pi d * \Sigma_i^{(m)} = +2\pi * j_i^{(m)} \neq 0, \quad (7)$$

where $d^2 B_i = 0$ and $d * \Sigma_i^{(m)} = - * j_i^{(m)}$ are used. Since the latter relation is also written as

$$\delta \Sigma_i^{(m)} = -j_i^{(m)}, \quad (8)$$

one finds that the color-electric current $j_i^{(m)}$ is defined as the boundary of the Dirac string $\Sigma_i^{(m)}$. In order to treat the Dirac string singularity further, it is useful to decompose the dual gauge field B_i into two parts called regular and singular ones,

$$B_i = B_i^{\text{reg}} + B_i^{\text{sing}}, \quad (9)$$

where the singular part is explicitly written as

$$B_i^{\text{sing}} = 2\pi \Delta^{-1} \delta * \Sigma_i^{(m)}. \quad (10)$$

The inverse of the Laplacian, Δ^{-1} , is the Coulomb propagator ($\Delta \Delta^{-1} = \Delta^{-1} \Delta = 1$). Note that Δ and Δ^{-1} commute with the exterior derivative d , the codifferential δ , and the Hodge star $*$. Notice further that B_i^{sing} is a background gauge field associated with the singular string surface. By using the relation $d\Delta\delta + \delta\Delta d = 1$, one finds that Eq. (10) leads to

$$dB_i^{\text{sing}} = 2\pi (*\Sigma_i^{(m)} + *C_i^{(m)}). \quad (11)$$

Then, the dual field strength is written as

$$F_i = dB_i^{\text{reg}} + 2\pi * C_i^{(m)}, \quad (12)$$

where

$$*C_i^{(m)} = \Delta^{-1} \delta * j_i^{(m)}. \quad (13)$$

Clearly, from this expression we see the violation of the dual Bianchi identity, since $d^2 B_i^{\text{reg}} = 0$ and $d * C_i^{(m)} = * j_i^{(m)}$. Then, the DGL action is written as

$$S_{\text{DGL}} = \sum_{i=1}^3 \left[\frac{\beta_g}{2} \left(dB_i^{\text{reg}} + 2\pi * C_i^{(m)} \right)^2 + (d\phi_i)^2 + ((B_i^{\text{reg}} + B_i^{\text{sing}} + d\eta_i^{\text{reg}})\phi_i)^2 + \lambda(\phi_i^2 - v^2)^2 \right], \quad (14)$$

where the polar decomposition form of the monopole field $\chi_i = \phi_i \exp(i\eta_i)$ ($\phi_i, \eta_i \in \mathbb{R}$) is inserted. Since we are interested in the *open*-string system, we investigate the case that the phase of the monopole field is single-valued, so that we have $\eta = \eta^{\text{reg}}$. In general the phase η can contain the singular part, which describes a *closed*-string, corresponding physically to the glueball system [28]. Such contribution should be taken into account when glueballs and quantum effects in the dual superconducting vacuum are concerned.

4 Functional path integration and the effective string action

The effective string action $S_{\text{eff}}(\Sigma_i^{(m)})$, written in terms of the color-magnetic representation of the color-electric Dirac string, will be obtained after the

functional path-integration over both the dual gauge fields and the monopole fields. Explicitly, this can be expressed by the following relation:

$$\begin{aligned}\mathcal{Z} &= \int \left(\prod_{i=1}^3 \mathcal{D}B_i^{\text{reg}} \delta[\delta B_i^{\text{reg}} - f_i] \right) \left(\prod_{i=1}^3 \mathcal{D}\phi_i^2 \mathcal{D}\eta_i^{\text{reg}} \right) \delta\left[\sum_{i=1}^3 \delta B_i^{\text{reg}}\right] \\ &\quad \times \exp[-S_{\text{DGL}}(B_i^{\text{reg}}, \phi_i, \eta_i^{\text{reg}}, \Sigma_i^{(m)})] \\ &= \exp[-S_{\text{eff}}(\Sigma_i^{(m)})].\end{aligned}\tag{15}$$

It is important to note that, when we adopt the Weyl symmetric formulation, we need to take into account the (Lorentz invariant) δ -function constraint among the dual gauge fields as

$$\sum_{i=1}^3 \delta B_i^{\text{reg}} = 0.\tag{16}$$

As a result, the dual gauge symmetry of the DGL theory is kept in $U(1) \times U(1)$. In fact, although the DGL action (1) itself has the $[U(1)]^3$ dual gauge symmetry

$$\chi_i \mapsto \chi_i \exp(i\alpha_i), \quad B_i^{\text{reg}} \mapsto B_i^{\text{reg}} - d\alpha_i \quad (i = 1, 2, 3),\tag{17}$$

this is finally reduced to $U(1) \times U(1)$ by the relation $\sum_{i=1}^3 d\alpha_i = 0$, which is implied by the constraint (16).²

Let us perform next the functional path integration over fields. For this aim, the quadratic term of the dual field strength in the action is evaluated as

$$\frac{\beta_g}{2} \left(dB_i^{\text{reg}} + 2\pi * C_i^{(m)} \right)^2 = \frac{\beta_g}{2} (dB_i^{\text{reg}})^2 + 2\pi^2 \beta_g \left(j_i^{(m)}, \Delta^{-1} j_i^{(m)} \right),\tag{18}$$

where we use

$$(*C_i^{(m)}, *C_i^{(m)}) = \left(j_i^{(m)}, \Delta^{-1} j_i^{(m)} \right),\tag{19}$$

and the cross term vanishes. Next, inserting an identity for auxiliary fields E_i (1-form, $i = 1, 2, 3$),

² Here it may be interesting to comment that in this formulation we do not need an additional constraint for the phase of the monopole field such as $\sum_{i=1}^3 \eta_i = \text{const.}$ In fact, such a requirement is already implicitly taken into account by Eq. (16). For instance, the special choice of the gauge, (unitary gauge) $\alpha_i \equiv -\eta_i$ ($i = 1, 2, 3$), leads to the corresponding constraint among the phase of the monopole field.

$$\begin{aligned}
& \exp \left[-((B_i^{\text{reg}} + B_i^{\text{sing}} + d\eta_i^{\text{reg}})\phi_i)^2 \right] \\
& = \phi_i^{-4} \int \mathcal{D}E_i \exp \left[- \left\{ \frac{1}{4}(E_i, \frac{1}{\phi_i^2}E_i) - i(B_i^{\text{reg}} + B_i^{\text{sing}} + d\eta_i^{\text{reg}}, E_i) \right\} \right], \quad (20)
\end{aligned}$$

where, in general, $\sum_{i=1}^3 E_i \neq 0$, and rewriting the δ -function constraint for the dual gauge fields as a Fourier integral, the partition function takes the form

$$\begin{aligned}
\mathcal{Z} = & \int \mathcal{D}k \left(\prod_{i=1}^3 \mathcal{D}B_i^{\text{reg}} \delta[\delta B_i^{\text{reg}} - f_{B\,i}] \right) \left(\prod_{i=1}^3 \mathcal{D}\phi_i^{-2} \mathcal{D}\eta_i^{\text{reg}} \right) \left(\prod_{i=1}^3 \mathcal{D}E_i \right) \\
& \times \exp \left[- \sum_{i=1}^3 \left\{ 2\pi^2 \beta_g \left(j_i^{(m)}, \Delta^{-1} j_i^{(m)} \right) + \frac{\beta_g}{2} (dB_i^{\text{reg}})^2 + (d\phi_i)^2 + \frac{1}{4} (E_i, \frac{1}{\phi_i^2} E_i) \right. \right. \\
& \quad \left. \left. - i(B_i^{\text{sing}}, E_i) - i(B_i^{\text{reg}}, E_i + dk) - i(d\eta_i^{\text{reg}}, E_i) + \lambda(\phi_i^2 - v^2)^2 \right\} \right]. \quad (21)
\end{aligned}$$

Here, k is a 1-form auxiliary field. Notice that, due to $(d\eta_i^{\text{reg}}, E_i) = (\eta_i^{\text{reg}}, \delta E_i)$, the integrations over the phase η_i^{reg} ($i = 1, 2, 3$) lead to the constraint $\delta E_i = 0$ which is resolved by introducing the Kalb-Ramond (KR) fields h_i (2-form, $i = 1, 2, 3$) as

$$\delta[\delta E_i] = \int \mathcal{D}h_i \delta[\delta h_i - f_{h\,i}] \delta[E_i - \delta * h_i], \quad (22)$$

where, in general, $\sum_{i=1}^3 h_i \neq 0$. Due to the hyper-gauge invariance $h \mapsto h + d\Lambda$ with a 1-form field Λ , the hyper-gauge fixing functional also appears in order to avoid the overcounting in the integration over h . Then, we have

$$\begin{aligned}
\mathcal{Z} = & \int \mathcal{D}k \left(\prod_{i=1}^3 \mathcal{D}B_i^{\text{reg}} \delta[\delta B_i^{\text{reg}} - f_{B\,i}] \right) \left(\prod_{i=1}^3 \mathcal{D}\phi_i^{-2} \right) \\
& \times \left(\prod_{i=1}^3 \mathcal{D}E_i \mathcal{D}h_i \delta[\delta h_i - f_{h\,i}] \delta[E_i - \delta * h_i] \right) \\
& \times \exp \left[- \sum_{i=1}^3 \left\{ 2\pi^2 \beta_g \left(j_i^{(m)}, \Delta^{-1} j_i^{(m)} \right) + \frac{\beta_g}{2} (dB_i^{\text{reg}})^2 + (d\phi_i)^2 \right. \right. \\
& \quad \left. \left. + \frac{1}{4} (E_i, \frac{1}{\phi_i^2} E_i) - i(B_i^{\text{sing}}, E_i) - i(B_i^{\text{reg}}, E_i + dk) + \lambda(\phi_i^2 - v^2)^2 \right\} \right]. \quad (23)
\end{aligned}$$

Now the integration over E_i ($i = 1, 2, 3$) leads to

$$\begin{aligned}
\mathcal{Z} = & \int \mathcal{D}k \left(\prod_{i=1}^3 \mathcal{D}B_i^{\text{reg}} \delta[\delta B_i^{\text{reg}} - f_{B\,i}] \right) \left(\prod_{i=1}^3 \mathcal{D}\phi_i^{-2} \right) \left(\prod_{i=1}^3 \mathcal{D}h_i \delta[\delta h_i - f_{h\,i}] \right) \\
& \times \exp \left[- \sum_{i=1}^3 \left\{ 2\pi^2 \beta_g \left(j_i^{(m)}, \Delta^{-1} j_i^{(m)} \right) + \frac{\beta_g}{2} (dB_i^{\text{reg}})^2 + (d\phi_i)^2 \right. \right. \\
& + \frac{1}{4} (\delta * h_i, \frac{1}{\phi_i^2} \delta * h_i) - i(B_i^{\text{sing}}, \delta * h_i) - i(B_i^{\text{reg}}, \delta * h_i + dk) \\
& \left. \left. + \lambda(\phi_i^2 - v^2)^2 \right\} \right]. \tag{24}
\end{aligned}$$

In order to evaluate the partition function further, we use the relations

$$\begin{aligned}
\sum_{i=1}^3 (B_i^{\text{sing}}, \delta * h_i) &= 2\pi \sum_{i=1}^3 (\Delta^{-1} \delta * \Sigma_i^{(m)}, \delta * h_i) \\
&= 2\pi \sum_{i=1}^3 (h_i, \Sigma_i^{(m)}) + 2\pi \sum_{i=1}^3 (\delta h_i, \Delta^{-1} j_i^{(m)}), \tag{25}
\end{aligned}$$

and

$$(\delta * h_i, \frac{1}{\phi_i^2} \delta * h_i) = (dh_i, \frac{1}{\phi_i^2} dh_i). \tag{26}$$

By inserting an identity in form of a path-integration over $f_{B\,i}$,

$$const. = \int \mathcal{D}f_{B\,i} \exp \left[- \frac{\beta_g}{2\xi_{B\,i}} f_{B\,i}^2 \right], \tag{27}$$

and taking $\xi_{B\,i} = 1$, we can integrate over B_i^{reg} as

$$\begin{aligned}
\mathcal{Z} = & \int \mathcal{D}k \left(\prod_{i=1}^3 \mathcal{D}\phi_i^{-2} \mathcal{D}h_i \delta[\delta h_i - f_{h\,i}] \right) \\
& \times \exp \left[- \sum_{i=1}^3 \left\{ 2\pi^2 \beta_g \left(j_i^{(m)}, \Delta^{-1} j_i^{(m)} \right) \right. \right. \\
& + \frac{1}{2\beta_g} (\delta * h_i + dk, \Delta^{-1} (\delta * h_i + dk)) + (d\phi_i)^2 + \frac{1}{4} (dh_i, \frac{1}{\phi_i^2} dh_i) \\
& \left. \left. - 2\pi i (h_i, \Sigma_i^{(m)}) - 2\pi i (\delta h_i, \Delta^{-1} j_i^{(m)}) + \lambda(\phi_i^2 - v^2)^2 \right\} \right], \tag{28}
\end{aligned}$$

where

$$(\delta * h_i + dk, \Delta^{-1} (\delta * h_i + dk)) = (h_i)^2 - (\delta h_i, \Delta^{-1} \delta h_i) + (k)^2. \tag{29}$$

Finally, the integration of k leads to a constant providing us with the expression

$$\begin{aligned} \mathcal{Z} = & \int \left(\prod_{i=1}^3 \mathcal{D}\phi_i^{-2} \mathcal{D}h_i \delta[\delta h_i - f_{hi}] \right) \\ & \times \exp \left[- \sum_{i=1}^3 \left\{ 2\pi^2 \beta_g (j_i^{(m)}, \Delta^{-1} j_i^{(m)}) + \frac{1}{2\beta_g} (h_i)^2 - \frac{1}{2\beta_g} (\delta h_i, \Delta^{-1} \delta h_i) \right. \right. \\ & + (d\phi_i)^2 + \frac{1}{4} (dh_i, \frac{1}{\phi_i^2} dh_i) - 2\pi i (h_i, \Sigma_i^{(m)}) - 2\pi i (\delta h_i, \Delta^{-1} j_i^{(m)}) \\ & \left. \left. + \lambda (\phi_i^2 - v^2)^2 \right\} \right]. \end{aligned} \quad (30)$$

As in our previous paper [39], we divide the action into three parts as

$$S = \sum_{i=1}^3 [S_i^{(1)} + S_i^{(2)} + S_i^{(3)}], \quad (31)$$

where

$$S_i^{(1)} = 2\pi^2 \beta_g (j_i^{(m)}, \Delta^{-1} j_i^{(m)}), \quad (32)$$

$$S_i^{(2)} = (d\phi_i)^2 + \frac{1}{4} (dh_i, \left\{ \frac{1}{\phi_i^2} - \frac{1}{v^2} \right\} dh_i) + \lambda (\phi_i^2 - v^2)^2, \quad (33)$$

$$\begin{aligned} S_i^{(3)} = & \frac{1}{4v^2} (dh_i)^2 + \frac{1}{2\beta_g} (h_i)^2 - \frac{1}{2\beta_g} (\delta h_i, \Delta^{-1} \delta h_i) \\ & - 2\pi i (h_i, \Sigma_i^{(m)}) - 2\pi i (\delta h_i, \Delta^{-1} j_i^{(m)}). \end{aligned} \quad (34)$$

Since these three actions have completely the same structures as in the SU(2) case [39] except for the labels of i , the further evaluation of these actions is straightforward. The first action $S_i^{(1)}$ provides the pure Coulombic potential for the static quark-antiquark system, since the color-electric currents interact via the Coulomb propagator. The second and third actions, $S_i^{(2)}$ and $S_i^{(3)}$, should be evaluated by taking into account the variation of the monopole field ϕ_i in the vicinity of the center of the flux tube, which defines the “core” region of the flux tube. In fact, its classical profile shows the behavior $\phi_i = v$ at the distant region from the center, while it tends to vanish as $\phi_i \rightarrow 0$ on the Dirac string if $\Sigma_i^{(m)} \neq 0$ (if $\Sigma_i^{(m)} = 0$, the monopole field behaves as $\phi_i = v$ everywhere [29]). Effectively, the core region can be characterized by the inverse of the monopole mass as $\rho \leq m_\chi^{-1}$. It is interesting to note that the second action contributes only inside of the core since it vanishes in the region where $\phi_i = v$, and its leading term contains the NG action with the

string tension σ^{core} [40]. At the same time, the third action in the core region contributes a boundary contribution $S_i^{\text{core}}(j_i^{(m)})$. Then, the action for the core region is

$$S_{i|<m_\chi^{-1}} = S_i^{\text{core}}(j_i^{(m)}) + \sigma^{\text{core}} \int_{\Sigma_i^{(m)}} d^2\xi_i \sqrt{g(\xi_i)}, \quad (35)$$

where ξ_i^a ($a = 1, 2$) parametrize the string world sheet $\tilde{x}_{i\mu}(\xi_i)$, and $g(\xi_i)$ is the determinant of the induced metric, $g_{ab}(\xi_i) \equiv \frac{\partial \tilde{x}_{i\mu}(\xi_i)}{\partial \xi_i^a} \frac{\partial \tilde{x}_{i\mu}(\xi_i)}{\partial \xi_i^b}$ [40,41]. Here, the third action in the outside region $\rho > m_\chi^{-1}$ is responsible for the description of the “surface” of the flux tube. By inserting a suitable identity in form of a path-integral over the hyper-gauge fixing function f_{hi} ,

$$\begin{aligned} \text{const.} &= \int \mathcal{D}f_{hi} \\ &\times \exp \left[-\frac{1}{4v^2} \left([f_{hi} + 4\pi v^2 i D j_i^{(m)}], D^{-1} \Delta^{-1} [f_{hi} + 4\pi v^2 i D j_i^{(m)}] \right) \right], \end{aligned} \quad (36)$$

where $D \equiv (\Delta + m_B^2)^{-1}$ is the propagator of the massive KR field, we can further integrate over the KR field as

$$\begin{aligned} S_{i|>m_\chi^{-1}} &= 4\pi^2 v^2 \left(\Sigma_i^{(m)}, D \Sigma_i^{(m)} \right) \Big|_{>m_\chi^{-1}} \\ &+ 2\pi^2 \beta_g \left(j_i^{(m)}, [D - \Delta^{-1}] j_i^{(m)} \right) \Big|_{>m_\chi^{-1}}. \end{aligned} \quad (37)$$

Now “ $|_{>m_\chi^{-1}}$ ” means that a corresponding effective cutoff should be taken into account. One finds that the first term describes the interaction between the color-electric Dirac string via the propagator of the massive KR field. The derivative expansion of the KR propagator with respect to the covariant Laplacian on the string world sheet which is defined by

$$\Delta_{\xi_i} = -\frac{1}{\sqrt{g(\xi_i)}} \partial_a g^{ab}(\xi_i) \sqrt{g(\xi_i)} \partial_b, \quad (38)$$

leads to the following form:

$$\begin{aligned}
& 4\pi^2 v^2 \left(\Sigma_i^{(m)}, D\Sigma_i^{(m)} \right) \Big|_{>m_\chi^{-1}} \\
&= \sigma^{\text{surf}} \int_{\Sigma_i^{(m)}} d^2 \xi_i \sqrt{g(\xi_i)} \\
&+ \alpha^{\text{surf}} \int_{\Sigma_i^{(m)}} d^2 \xi_i \sqrt{g(\xi_i)} g^{ab}(\xi_i) (\partial_a t_{\mu\nu}(\xi_i)) (\partial_b t_{\mu\nu}(\xi_i)) + \mathcal{O}(\Delta_{\xi_i}^2), \quad (39)
\end{aligned}$$

where $t_{\mu\nu}(\xi_i) = \frac{\epsilon^{ab}}{\sqrt{g(\xi_i)}} \frac{\partial \tilde{x}_\mu(\xi_i)}{\partial \xi_i^a} \frac{\partial \tilde{x}_\nu(\xi_i)}{\partial \xi_i^b}$ is an antisymmetric tensor which determines the orientation of the string world sheet $\Sigma_i^{(m)}$. The first term represents the NG action with the string tension

$$\sigma^{\text{surf}} = \pi v^2 \ln \frac{m_B^2 + m_\chi^2}{m_B^2} = \pi v^2 \ln (1 + \kappa^2), \quad (40)$$

and the second term is the so-called rigidity term with the negative coefficient

$$\alpha^{\text{surf}} = \frac{\pi v^2}{2} \left(\frac{1}{m_\chi^2 + m_B^2} - \frac{1}{m_B^2} \right) = -\frac{\pi \beta_g}{4} \frac{\kappa^2}{1 + \kappa^2} \quad (< 0), \quad (41)$$

where $\kappa = m_\chi/m_B$ is the GL parameter.

Finally, by combining all contributions, the pure Coulomb term (32), the core term (35), and the surface term (37) with derivative expansion (39), we get the effective string action

$$\begin{aligned}
S_{\text{eff}} = & \sum_{i=1}^3 \left[S_i(j_i^{(m)}) + (\sigma^{\text{core}} + \sigma^{\text{surf}}) \int_{\Sigma_i^{(m)}} d^2 \xi_i \sqrt{g(\xi_i)} \right. \\
& \left. + \alpha^{\text{surf}} \int_{\Sigma_i^{(m)}} d^2 \xi_i \sqrt{g(\xi_i)} g^{ab}(\xi_i) (\partial_a t_{\mu\nu}(\xi_i)) (\partial_b t_{\mu\nu}(\xi_i)) + \mathcal{O}(\Delta_{\xi_i}^2) \right], \quad (42)
\end{aligned}$$

where the boundary (color-electric current) contribution is given by

$$\begin{aligned}
S_i(j_i^{(m)}) = & S_i^{\text{core}}(j_i^{(m)}) + 2\pi^2 \beta_g(j_i^{(m)}, \Delta^{-1} j_i^{(m)}) \\
& + 2\pi^2 \beta_g(j_i^{(m)}, [D - \Delta^{-1}] j_i^{(m)}) \Big|_{>m_\chi^{-1}}. \quad (43)
\end{aligned}$$

Eqs. (42), (43) are the main results of this paper .

It has to be noted that the boundary term can be simplified only in the London limit $m_\chi \rightarrow \infty$. In this limit, the first term disappears, $S_i^{\text{core}}(j_i^{(m)}) \rightarrow 0$, and




Fig. 1. The structure of the color-electric Dirac string in the color-magnetic representation for the mesonic (left) and the baryonic (right) quark systems.

the third term becomes an “exact” Yukawa minus Coulomb term. Then, the *cancellation of the Coulomb term takes place*, and finally the boundary term is reduced into the ordinary Yukawa interaction as

$$S_i(j_i^{(m)}) \rightarrow 2\pi^2\beta_g(j_i^{(m)}, Dj_i^{(m)}). \quad (44)$$

However, if the vacuum is not in the deep type-II limit, the boundary contribution has, in principle, a complicated structure which can only be investigated numerically.

Clearly, the explicit form of the effective string action depends on the quark system, that is to say, the combination of the color-electric charges. In the SU(3) case, one can consider not only the mesonic string, described by $R\text{-}\bar{R}$, $B\text{-}\bar{B}$ and $G\text{-}\bar{G}$ systems, but also the *baryonic* string, as given by the combination of three color-electric charges $R\text{-}B\text{-}G$ (see, Fig. 1). Remarkably, since the form of the action is now the Weyl symmetric one, which means invariant under the exchange of the color labels i , one can immediately conclude that all mesonic strings do not depend on the color. The baryonic system is constructed by connecting the three-colored mesonic strings, $R\text{-}\bar{R}$, $B\text{-}\bar{B}$, and $G\text{-}\bar{G}$, at a certain *junction*, forming a Y-shaped structure. This is possible because the sum of all three color charges at the same point turns out to be zero. In Ref. [37], the static potential among three ends is calculated using the effective string action in the London limit and its short distance region is described by the Yukawa potential, which is consistent with our estimation in Eq. (44). The profile of the baryonic flux-tube solution in the DGL theory is investigated in Refs. [27,30], where also the role of the color-electric Dirac string inside the flux tube is clarified.

5 Summary

In this paper we have studied the effective string action of the color-electric flux tube in the dual Ginzburg-Landau (DGL) theory using the path-integral analysis in differential forms, thus extending our previous work for the $U(1)$ dual Abelian Higgs (DAH) model [39] to the $U(1) \times U(1)$ case. By adopting the Weyl symmetric formulation, where the $U(1) \times U(1)$ symmetry is naturally extended to $[U(1)]^3$, the treatment of the DGL theory is found to be rather simple, since the resulting form is the sum of three types of the $U(1)$ DAH model. Taking advantage of this feature, we could similarly get the effective string action, which is expressed as the sum of the string action in the $U(1)$ case. Then the interpretation of the action is also straightforward. The difference is only the number of color-electric charges, which then allows us to describe not only the mesonic system but also the baryonic system. The latter is a typical system in the $SU(3)$ gluodynamics. In the present paper, we do not describe the baryon systems explicitly, which will be investigated in a future analysis.

Let us make some comments concerning the form of the effective string action in Eqs. (42) and (43). Analogously to the case of $SU(2)$, the effect of the finite thickness of the flux tube leads us again to a modified Yukawa interaction as the boundary contribution of the open string, which turns over to the ordinary Yukawa one only in the London limit. The derivative expansion of the KR propagator with respect to the covariant Laplacian (38) leads to the NG action and the rigidity term as the leading and next-to-leading parts of the action. Such an expansion is clearly justified when the fluctuation of the string world sheet can be treated perturbatively. If not, one has to handle the expression of the first term in Eq. (37) directly.

Finally, it is interesting to note that we can refer to the interaction among string world sheets of various colors without solving their dynamics. Since the sum of two types of color charges reduces to another type of *anti*-color charge, for instance as $R + B \rightarrow \bar{G}$, one finds that if two strings composed of R - $\bar{R}$ and B - $\bar{B}$ interact with each other, this will be equivalent to a $\bar{G}$ - G string. Then, it is easy to imagine that the force between them must be attractive.

One of the authors (Y.K.) is partially supported by the Ministry of Education, Science, Sports and Culture, Japan, Grant-in-Aid for Encouragement of Young Scientists (B), 14740161, 2002. We would like to thank D. Antonov and E.-M. Ilgenfritz for useful discussions.

References

- [1] G. 't Hooft, in *High-Energy Physics. Proceedings of the EPS International Conference, Palermo, Italy, 23-28 June 1975, Vol. 2*, edited by A. Zichichi, pp. 1225–1249, Bologna, 1976.
- [2] S. Mandelstam, Phys. Rept. **23C**, 245 (1976).
- [3] A. A. Abrikosov, Sov. Phys. JETP **5**, 1174 (1957).
- [4] H. B. Nielsen and P. Olesen, Nucl. Phys. **B61**, 45 (1973).
- [5] Y. Nambu, Phys. Rev. **D10**, 4262 (1974).
- [6] A. S. Kronfeld, G. Schierholz, and U. J. Wiese, Nucl. Phys. **B293**, 461 (1987).
- [7] H. Shiba and T. Suzuki, Phys. Lett. **B343**, 315 (1995), hep-lat/9406010.
- [8] H. Shiba and T. Suzuki, Phys. Lett. **B351**, 519 (1995), hep-lat/9408004.
- [9] Z. F. Ezawa and A. Iwazaki, Phys. Rev. **D25**, 2681 (1982).
- [10] Z. F. Ezawa and A. Iwazaki, Phys. Rev. **D26**, 631 (1982).
- [11] T. Suzuki and I. Yotsuyanagi, Phys. Rev. **D42**, 4257 (1990).
- [12] F. Brandstaeter, U. J. Wiese, and G. Schierholz, Phys. Lett. **B272**, 319 (1991).
- [13] O. Miyamura, Nucl. Phys. Proc. Suppl. **42**, 538 (1995).
- [14] J. D. Stack, S. D. Neiman, and R. J. Wensley, Phys. Rev. **D50**, 3399 (1994), hep-lat/9404014.
- [15] O. Miyamura, Phys. Lett. **B353**, 91 (1995).
- [16] G. 't Hooft, Nucl. Phys. **B190**, 455 (1981).
- [17] S. Maedan and T. Suzuki, Prog. Theor. Phys. **81**, 229 (1989).
- [18] T. Suzuki, Prog. Theor. Phys. **80**, 929 (1988).
- [19] H. Suganuma, S. Sasaki, and H. Toki, Nucl. Phys. **B435**, 207 (1995), hep-ph/9312350.
- [20] S. Sasaki, H. Suganuma, and H. Toki, Prog. Theor. Phys. **94**, 373 (1995).
- [21] V. Singh, D. A. Browne, and R. W. Haymaker, Phys. Lett. **B306**, 115 (1993), hep-lat/9301004.
- [22] Y.-c. Peng and R. W. Haymaker, Phys. Rev. **D52**, 3030 (1995), hep-lat/9503015.
- [23] Y. Matsubara, S. Ejiri, and T. Suzuki, Nucl. Phys. Proc. Suppl. **34**, 176 (1994), hep-lat/9311061.

- [24] K. Schilling, G. S. Bali, and C. Schlichter, Nucl. Phys. Proc. Suppl. **73**, 638 (1999), hep-lat/9809039.
- [25] F. V. Gubarev, E.-M. Ilgenfritz, M. I. Polikarpov, and T. Suzuki, Phys. Lett. **B468**, 134 (1999), hep-lat/9909099.
- [26] V. Bornyakov *et al.*, Nucl. Phys. Proc. Suppl. **106**, 634 (2002), hep-lat/0111042.
- [27] S. Kamizawa, Y. Matsubara, H. Shiba, and T. Suzuki, Nucl. Phys. **B389**, 563 (1993).
- [28] Y. Koma, H. Suganuma, and H. Toki, Phys. Rev. **D60**, 074024 (1999), hep-ph/9902441.
- [29] Y. Koma and H. Toki, Phys. Rev. **D62**, 054027 (2000), hep-ph/0004177.
- [30] Y. Koma, E.-M. Ilgenfritz, T. Suzuki, and H. Toki, Phys. Rev. **D64**, 014015 (2001), hep-ph/0011165.
- [31] P. Orland, Nucl. Phys. **B428**, 221 (1994), hep-th/9404140.
- [32] M. Sato and S. Yahikozawa, Nucl. Phys. **B436**, 100 (1995), hep-th/9406208.
- [33] E. T. Akhmedov, M. N. Chernodub, M. I. Polikarpov, and M. A. Zubkov, Phys. Rev. **D53**, 2087 (1996), hep-th/9505070.
- [34] D. Antonov and D. Ebert, Eur. Phys. J. **C8**, 343 (1999), hep-th/9806153.
- [35] A. M. Polyakov, Nucl. Phys. **B268**, 406 (1986).
- [36] H. Kleinert, Phys. Lett. **B174**, 335 (1986).
- [37] M. N. Chernodub and D. A. Komarov, JETP Lett. **68**, 117 (1998), hep-th/9809183.
- [38] D. Antonov and D. Ebert, Phys. Lett. **B444**, 208 (1998), hep-th/9809018.
- [39] Y. Koma, M. Koma, D. Ebert, and H. Toki, JHEP **08**, 047 (2002), hep-th/0108138.
- [40] D. Foerster, Nucl. Phys. **B81**, 84 (1974).
- [41] R. L. Davis and E. P. S. Shellard, Phys. Lett. **B214**, 219 (1988).